

# The maximal value of Dirichlet kernels with respect to representative product systems

R. Toledo

College of Nyíregyháza

September 21th, 2009  
Budapest

# The complete product of finite groups

We deal with Fourier analysis on a generalization of **Walsh-Paley** and **Vilenkin** systems.

The group  $\left( G := \prod_{k=0}^{\infty} G_k \right)$

Denote by  $G$  the **complete product of arbitrary finite groups** of order  $m_k$  ( $m_k \geq 2$ ,  $k \in \mathbf{N}$ ), with discrete topology and assign each singleton the measure  $\frac{1}{m_k}$ .  $G$  has the product topology and measure. (Haar measure)

# The complete product of finite groups

We deal with Fourier analysis on a generalization of **Walsh-Paley** and **Vilenkin** systems.

The group  $\left( G := \prod_{k=0}^{\infty} G_k \right)$

Denote by  $G$  the **complete product of arbitrary finite groups** of order  $m_k$  ( $m_k \geq 2$ ,  $k \in \mathbf{N}$ ), with discrete topology and assign each singleton the measure  $\frac{1}{m_k}$ .  $G$  has the product topology and measure. (Haar measure)

The dual object  $(\Sigma_k)$  of the finite group  $G_k$  ( $k \in \mathbf{N}$ )

is the set of all continuous irreducible unitary representations of the group  $G_k$  which are not equivalents.

The Coordinate functions

For any  $\sigma \in \Sigma_k$ , let  $\{\xi_1, \dots, \xi_{d_\sigma}\}$  be a fixed basis of the representation space of a representation  $U^{(\sigma)}$  in the class  $\sigma$  having the dimension  $d_\sigma$ .

The Coordinate functions:

$$u_{i,j}^{(\sigma)}(x) := \langle U_x^{(\sigma)} \xi_i, \xi_j \rangle, \quad i, j \in \{1, \dots, d_\sigma\}, \sigma \in \Sigma_k$$

The dual object  $(\Sigma_k)$  of the finite group  $G_k$  ( $k \in \mathbf{N}$ )

is the set of all continuous irreducible unitary representations of the group  $G_k$  which are not equivalents.

## The Coordinate functions

For any  $\sigma \in \Sigma_k$ , let  $\{\xi_1, \dots, \xi_{d_\sigma}\}$  be a fixed basis of the representation space of a representation  $U^{(\sigma)}$  in the class  $\sigma$  having the dimension  $d_\sigma$ .

The **Coordinate functions**:

$$u_{i,j}^{(\sigma)}(x) := \langle U_x^{(\sigma)} \xi_i, \xi_j \rangle, \quad i, j \in \{1, \dots, d_\sigma\}, \sigma \in \Sigma_k$$

# Orthonormal systems on finite groups

## The system $\varphi_k$

We order the all normalized coordinate functions of the finite group  $G_k$  ( $\varphi_k^0(x) = 1$ ) to obtain exactly  $m_k$  number of functions.

$$\varphi_k^s(x) = \sqrt{d_\sigma} u_{i,j}^{(\sigma)}(x) \quad (x \in G_k, s = 0, \dots, m_k - 1),$$

where  $\sigma \in \Sigma_k, i, j \in \{1, \dots, d_\sigma\}$ .

## Theorem

*The system  $\varphi_k$  is an orthonormal and complete system on  $L^2(G_k)$ .*

# Orthonormal systems on finite groups

## The system $\varphi_k$

We order the all normalized coordinate functions of the finite group  $G_k$  ( $\varphi_k^0(x) = 1$ ) to obtain exactly  $m_k$  number of functions.

$$\varphi_k^s(x) = \sqrt{d_\sigma} u_{i,j}^{(\sigma)}(x) \quad (x \in G_k, s = 0, \dots, m_k - 1),$$

where  $\sigma \in \Sigma_k, i, j \in \{1, \dots, d_\sigma\}$ .

## Theorem

*The system  $\varphi_k$  is an orthonormal and complete system on  $L^2(G_k)$ .*

# Example 1: Cyclic groups of order $m$ , $\mathbb{Z}_m$

## The generalized Rademacher functions

$$\varphi^s(x) = \exp(2\pi i s x / m) \quad (s \in \{0, \dots, m-1\}, x \in \mathbb{Z}_m, i^2 = -1)$$

- All of the members of system  $\varphi$  are characters.
- $|\varphi^s(x)| = 1$  for all  $x \in \mathbb{Z}_m$  and  $s \in \{0, \dots, m-1\}$ .
- $\|\varphi^s\|_1 = 1, \quad \|\varphi^s\|_\infty = 1$ .

# Example 1: Cyclic groups of order $m$ , $\mathbb{Z}_m$

The generalized Rademacher functions

$$\varphi^s(x) = \exp(2\pi i s x / m) \quad (s \in \{0, \dots, m-1\}, x \in \mathbb{Z}_m, i^2 = -1)$$

- All of the members of system  $\varphi$  are characters.
- $|\varphi^s(x)| = 1$  for all  $x \in \mathbb{Z}_m$  and  $s \in \{0, \dots, m-1\}$ .
- $\|\varphi^s\|_1 = 1, \quad \|\varphi^s\|_\infty = 1.$

# Example 1: Cyclic groups of order $m$ , $\mathbb{Z}_m$

## The generalized Rademacher functions

$$\varphi^s(x) = \exp(2\pi i s x / m) \quad (s \in \{0, \dots, m-1\}, x \in \mathbb{Z}_m, i^2 = -1)$$

- All of the members of system  $\varphi$  are characters.
- $|\varphi^s(x)| = 1$  for all  $x \in \mathbb{Z}_m$  and  $s \in \{0, \dots, m-1\}$ .
- $\|\varphi^s\|_1 = 1, \quad \|\varphi^s\|_\infty = 1.$

## The generalized Rademacher functions

$$\varphi^s(x) = \exp(2\pi i s x / m) \quad (s \in \{0, \dots, m-1\}, x \in \mathbb{Z}_m, i^2 = -1)$$

- All of the members of system  $\varphi$  are characters.
- $|\varphi^s(x)| = 1$  for all  $x \in \mathbb{Z}_m$  and  $s \in \{0, \dots, m-1\}$ .
- $\|\varphi^s\|_1 = 1, \quad \|\varphi^s\|_\infty = 1.$

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

• The system has 2 characters and a representation of dimension 2.

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

- The system has 2 characters and a representation of dimension 2.
- $|\varphi^s(x)| = 0$  for some  $x \in \mathcal{X}_m$  and  $0 \leq s < 6$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 6$  and

$$\max_{0 \leq s < 6} \|\varphi^s\|_1 \|\varphi^s\|_\infty = \frac{4}{3}$$

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

- The system has 2 characters and a representation of dimension 2.
- $|\varphi^s(x)| = 0$  for some  $x \in \mathcal{Z}_m$  and  $0 \leq s < 6$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 6$  and

$$\max_{s=0 \dots 5} \|\varphi^s\|_1 \|\varphi^s\|_\infty = \frac{4}{3}.$$

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

- The system has 2 characters and a representation of dimension 2.
- $|\varphi^s(x)| = 0$  for some  $x \in \mathcal{Z}_m$  and  $0 \leq s < 6$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 6$  and

$$\max_{s=0\dots 5} \|\varphi^s\|_1 \|\varphi^s\|_\infty = \frac{4}{3}.$$

# Example 2: The permutation group of 3 elements, $S_3$

	$e$	$(12)$	$(13)$	$(23)$	$(123)$	$(132)$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1
$\varphi^1$	1	-1	-1	-1	1	1	1	1
$\varphi^2$	$\sqrt{2}$	$-\sqrt{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^3$	$\sqrt{2}$	$\sqrt{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	$\frac{2\sqrt{2}}{3}$	$\sqrt{2}$
$\varphi^4$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$
$\varphi^5$	0	0	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$-\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{2}$	$\frac{\sqrt{6}}{3}$	$\frac{\sqrt{6}}{2}$

- The system has 2 characters and a representation of dimension 2.
- $|\varphi^s(x)| = 0$  for some  $x \in \mathcal{Z}_m$  and  $0 \leq s < 6$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 6$  and

$$\max_{s=0\dots 5} \|\varphi^s\|_1 \|\varphi^s\|_\infty = \frac{4}{3}.$$

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

- The system has 4 characters and a representation of dimension 2.
- Monomial  $\chi^s = \chi^t$  iff  $s \equiv t \pmod{4}$ .

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

- The system has 4 characters and a representation of dimension 2.
- Monomial:  $|\varphi^s(x)| = 0$  or  $|\varphi^s(x)| = \sqrt{d_s}$  for all  $x \in \mathcal{X}_m$  and  $0 \leq s < 8$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 8$  but

$$\max_{0 \leq s < 8} \|\varphi^s\|_1 \|\varphi^s\|_\infty = 1.$$

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

- The system has 4 characters and a representation of dimension 2.
- Monomial:  $|\varphi^s(x)| = 0$  or  $|\varphi^s(x)| = \sqrt{d_\sigma}$  for all  $x \in \mathcal{Z}_m$  and  $0 \leq s < 8$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 8$  but

$$\max_{s=0 \dots 7} \|\varphi^s\|_1 \|\varphi^s\|_\infty = 1.$$

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

- The system has 4 characters and a representation of dimension 2.
- Monomial:  $|\varphi^s(x)| = 0$  or  $|\varphi^s(x)| = \sqrt{d_\sigma}$  for all  $x \in \mathcal{Z}_m$  and  $0 \leq s < 8$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 8$  but

$$\max_{s=0 \dots 7} \|\varphi^s\|_1 \|\varphi^s\|_\infty = 1.$$

# Example 3: The quaternion group of order 8:

$$\mathcal{Q}_2 := \{[a, b] : a^4 = e, b^2 = a^2, bab^{-1} = a^3\}$$

	$e$	$a$	$a^2$	$a^3$	$b$	$ab$	$a^2b$	$a^3b$	$\ \varphi^s\ _1$	$\ \varphi^s\ _\infty$
$\varphi^0$	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	-1	-1	-1	-1	1	1
$\varphi^2$	1	-1	1	-1	1	-1	1	-1	1	1
$\varphi^3$	1	-1	1	-1	-1	1	-1	1	1	1
$\varphi^4$	$\sqrt{2}$	$\sqrt{2}i$	$-\sqrt{2}$	$-\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^5$	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	0	0	0	0	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^6$	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}i$	$-\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$
$\varphi^7$	0	0	0	0	$-\sqrt{2}$	$-\sqrt{2}i$	$\sqrt{2}$	$\sqrt{2}i$	$\frac{\sqrt{2}}{2}$	$\sqrt{2}$

- The system has 4 characters and a representation of dimension 2.
- **Monomial:**  $|\varphi^s(x)| = 0$  or  $|\varphi^s(x)| = \sqrt{d_\sigma}$  for all  $x \in \mathcal{Z}_m$  and  $0 \leq s < 8$ .
- $\|\varphi^s\|_\infty > 1$  for some  $0 \leq s < 8$  but

$$\max_{s=0 \dots 7} \|\varphi^s\|_1 \|\varphi^s\|_\infty = 1.$$

# Representative product systems

The  $m$ -adic expansion of  $n$ :  $(n_0, n_1, \dots)$

Denote  $M_0 := 1$  and  $M_{k+1} := m_k M_k$ , ( $k \in \mathbf{N}$ ). Given  $n \in \mathbf{N}$  it is possible to write  $n$  uniquely as

$$n = \sum_{k=0}^{\infty} n_k M_k, \quad (0 \leq n_k < m_k).$$

A representative product systems

G. Gát and R. Toledo

is the product system of  $\varphi$ :

$$\psi_n(x) := \prod_{k=0}^{\infty} \varphi_k^{n_k}(x_k) \quad (x \in G).$$

The  $m$ -adic expansion of  $n$ :  $(n_0, n_1, \dots)$

Denote  $M_0 := 1$  and  $M_{k+1} := m_k M_k$ , ( $k \in \mathbf{N}$ ). Given  $n \in \mathbf{N}$  it is possible to write  $n$  uniquely as

$$n = \sum_{k=0}^{\infty} n_k M_k, \quad (0 \leq n_k < m_k).$$

A representative product systems

G. Gát and R. Toledo

is the product system of  $\varphi$ :

$$\psi_n(x) := \prod_{k=0}^{\infty} \varphi_k^{n_k}(x_k) \quad (x \in G).$$

## Theorem

*A representative product system is an orthonormal and complete system on  $L^2(G)$ .*

Characteristics of the system  $\psi$  for noncommutative cases:

- It is not uniformly bounded.
- It takes the value 0.

## Theorem

*A representative product system is an orthonormal and complete system on  $L^2(G)$ .*

Characteristics of the system  $\psi$  for noncommutative cases:

- It is not uniformly bounded.
- It takes the value 0.

## Theorem

*A representative product system is an orthonormal and complete system on  $L^2(G)$ .*

Characteristics of the system  $\psi$  for noncommutative cases:

- It is not uniformly bounded.
- It takes the value 0.

# Convergence in $L^p$ -norm of Fourier series

The  $n$ -th partial sums of Fourier series

$$S_n f := \sum_{k=0}^{n-1} \widehat{f}_k \psi_k \quad (n \in \mathbf{N}), \quad \text{where } \widehat{f}_k := \int_G f \overline{\psi}_k d\mu.$$

Theorem (P. Simon, F. Schipp and W. S. Young)

*Let  $G$  be a Vilenkin group and  $1 < p < \infty$ . Then for all function  $f \in L^p(G)$  the sequence of partial sums  $S_n f$  of the Fourier series of  $f$  converges to the function  $f$  in  $L^p$ -norm.*

# Convergence in $L^p$ -norm of Fourier series

The  $n$ -th partial sums of Fourier series

$$S_n f := \sum_{k=0}^{n-1} \widehat{f}_k \psi_k \quad (n \in \mathbf{N}), \quad \text{where } \widehat{f}_k := \int_G f \overline{\psi_k} d\mu.$$

**Theorem (P. Simon, F. Schipp and W. S. Young)**

*Let  $G$  be a Vilenkin group and  $1 < p < \infty$ . Then for all function  $f \in L^p(G)$  the sequence of partial sums  $S_n f$  of the Fourier series of  $f$  converges to the function  $f$  in  $L^p$ -norm.*

# Convergence in $L^p$ -norm of Fourier series

The sequence  $\Psi$

$$\Psi_k = \prod_{i=0}^{k-1} \max_{s < m_i} \|\varphi_i^s\|_1 \|\varphi_i^s\|_\infty \quad (k \in \mathbf{N}).$$

Theorem (R. Toledo)

*If  $G$  is a bounded group with unbounded sequence  $\Psi$ , then for all  $p \neq 2$ ,  $1 < p < \infty$  there exists a function  $f \in L^p(G)$  such that the sequence of partial sums  $S_n f$  of the Fourier series of  $f$  does not converge to the function  $f$  in  $L^p$ -norm.*

The sequence  $\Psi$

$$\Psi_k = \prod_{i=0}^{k-1} \max_{s < m_i} \|\varphi_i^s\|_1 \|\varphi_i^s\|_\infty \quad (k \in \mathbf{N}).$$

## Theorem (R. Toledo)

*If  $G$  is a bounded group with unbounded sequence  $\Psi$ , then for all  $p \neq 2$ ,  $1 < p < \infty$  there exists a function  $f \in L^p(G)$  such that the sequence of partial sums  $S_n f$  of the Fourier series of  $f$  does not converge to the function  $f$  in  $L^p$ -norm.*

## The Dirichlet kernels

$$D_n(x, y) := \sum_{k=0}^{n-1} \psi_k(x) \overline{\psi_k(y)} \quad (n \in \mathbf{P})$$

$$S_n f(x) = \int_G f(y) D_n(x, y) d\mu(y)$$

## The maximal value of Dirichlet kernels

$$D_n := \sup_{x, y \in G} |D_n(x, y)| \quad (n \in \mathbf{P})$$

## The Dirichlet kernels

$$D_n(x, y) := \sum_{k=0}^{n-1} \psi_k(x) \overline{\psi_k(y)} \quad (n \in \mathbf{P})$$

$$S_n f(x) = \int_G f(y) D_n(x, y) d\mu(y)$$

## The maximal value of Dirichlet kernels

$$D_n := \sup_{x, y \in G} |D_n(x, y)| \quad (n \in \mathbf{P})$$

## The Dirichlet kernels

$$D_n(x, y) := \sum_{k=0}^{n-1} \psi_k(x) \overline{\psi_k(y)} \quad (n \in \mathbf{P})$$

$$S_n f(x) = \int_G f(y) D_n(x, y) d\mu(y)$$

## The maximal value of Dirichlet kernels

$$D_n := \sup_{x, y \in G} |D_n(x, y)| \quad (n \in \mathbf{P})$$

# The maximal value of Dirichlet kernels

If  $G$  is commutative then  $D_n = n$ , but for an arbitrary group  $G$

Theorem (R. Toledo)

If  $n \in \mathbf{P}$  and  $A := \max\{k \in \mathbf{N} : n_k \neq 0\}$ , then

$$n \leq D_n \leq M_{A+1}.$$


# The maximal value of Dirichlet kernels

If  $G$  is commutative then  $D_n = n$ , but for an arbitrary group  $G$

Theorem (R. Toledo)

If  $n \in \mathbf{P}$  and  $A := \max\{k \in \mathbf{N} : n_k \neq 0\}$ , then

$$n \leq D_n \leq M_{A+1}.$$



$D_n$  ( $n \leq 6^4$ ) on the complete product of  $\mathcal{S}_3$

Denote the indexes  $A$  and  $B$  by

$$A := \max\{k \in \mathbf{N} : n_k \neq 0\}, \quad \text{and} \quad B := \min\{k \in \mathbf{N} : n_k \neq 0, k \leq A\}.$$

It is not difficult to see that  $D_n = n$  if  $n$  satisfies the following two properties

- (i)  $\sum_{s=0}^{n_B-1} |\varphi_B^s(x_B)|^2 = n_B$  for all  $x_B \in G_B$ ,
- (ii)  $B = A$  or all of  $\varphi_i^{n_i}$  are characters for all  $B < i \leq A$  and  $x_i \in G_i$ .

$$D_n = n$$

The properties (i) and (ii) are not necessary for  $D_n = n$ . For instance, take the alternating group  $\mathcal{U}_4$  and  $\alpha = \exp(2\pi i/3)$ .

	e	(12)(34)	(13)(24)	(14)(23)	(123)	(142)	(134)	(234)	(124)	(132)	(234)	(143)
$\varphi^0$	1	1	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	$\alpha$	$\alpha$	$\alpha$	$\alpha$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$
$\varphi^2$	1	1	1	1	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha$	$\alpha$	$\alpha$	$\alpha$
$\varphi^3$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^4$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^5$	$\sqrt{3}$	$\sqrt{3}$	$-\sqrt{3}$	$-\sqrt{3}$	0	0	0	0	0	0	0	0
$\varphi^6$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^7$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^8$	0	0	0	0	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^9$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{10}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{11}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$

Take a new order:  $\{\varphi^0, \varphi^8, \varphi^5, \varphi^1, \varphi^2, \varphi^3, \varphi^4, \varphi^6, \varphi^7, \varphi^9, \varphi^{10}, \varphi^{11}\}$ .

Let  $G = \mathcal{Z}_2 \times \mathcal{U}_4 \times \mathcal{Z}_2 \times \mathcal{Z}_2 \times \dots$ , thus  $D_5 = 5$ , but property (ii) is not true because  $5 = (1, 2, 0, 0, \dots)$  and  $\varphi_1^2 = \varphi^5$  is not a character.

$$D_n = n$$

The properties (i) and (ii) are not necessary for  $D_n = n$ . For instance, take the alternating group  $\mathcal{U}_4$  and  $\alpha = \exp(2\pi i/3)$ .

	$e$	(12)(34)	(13)(24)	(14)(23)	(123)	(142)	(134)	(234)	(124)	(132)	(234)	(143)
$\varphi^0$	1	1	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	$\alpha$	$\alpha$	$\alpha$	$\alpha$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$
$\varphi^2$	1	1	1	1	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha$	$\alpha$	$\alpha$	$\alpha$
$\varphi^3$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^4$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^5$	$\sqrt{3}$	$\sqrt{3}$	$-\sqrt{3}$	$-\sqrt{3}$	0	0	0	0	0	0	0	0
$\varphi^6$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^7$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^8$	0	0	0	0	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^9$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{10}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{11}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$

Take a new order:  $\{\varphi^0, \varphi^8, \varphi^5, \varphi^1, \varphi^2, \varphi^3, \varphi^4, \varphi^6, \varphi^7, \varphi^9, \varphi^{10}, \varphi^{11}\}$ .

Let  $G = \mathcal{Z}_2 \times \mathcal{U}_4 \times \mathcal{Z}_2 \times \mathcal{Z}_2 \times \dots$ , thus  $D_5 = 5$ , but property (ii) is not true because  $5 = (1, 2, 0, 0, \dots)$  and  $\varphi_1^2 = \varphi^5$  is not a character.

$$D_n = n$$

The properties (i) and (ii) are not necessary for  $D_n = n$ . For instance, take the alternating group  $\mathcal{U}_4$  and  $\alpha = \exp(2\pi i/3)$ .

	$e$	(12)(34)	(13)(24)	(14)(23)	(123)	(142)	(134)	(234)	(124)	(132)	(234)	(143)
$\varphi^0$	1	1	1	1	1	1	1	1	1	1	1	1
$\varphi^1$	1	1	1	1	$\alpha$	$\alpha$	$\alpha$	$\alpha$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$
$\varphi^2$	1	1	1	1	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha^2$	$\alpha$	$\alpha$	$\alpha$	$\alpha$
$\varphi^3$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^4$	$\sqrt{3}$	$-\sqrt{3}$	0	0	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^5$	$\sqrt{3}$	$\sqrt{3}$	$-\sqrt{3}$	$-\sqrt{3}$	0	0	0	0	0	0	0	0
$\varphi^6$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^7$	0	0	$\sqrt{3}$	$-\sqrt{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$
$\varphi^8$	0	0	0	0	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^9$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{10}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$
$\varphi^{11}$	0	0	0	0	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{2}$

Take a new order:  $\{\varphi^0, \varphi^8, \varphi^5, \varphi^1, \varphi^2, \varphi^3, \varphi^4, \varphi^6, \varphi^7, \varphi^9, \varphi^{10}, \varphi^{11}\}$ .

Let  $G = \mathcal{Z}_2 \times \mathcal{U}_4 \times \mathcal{Z}_2 \times \mathcal{Z}_2 \times \dots$ , thus  $D_5 = 5$ , but property (ii) is not true because  $5 = (1, 2, 0, 0, \dots)$  and  $\varphi_1^2 = \varphi^5$  is not a character.

$$D_n = n$$

**Property (ii\*)**  $B = A$  or there exist  $r_i < d_i^{(n_i)}$  nonnegative integers such that

$$(d_i^{(n_i)} - 1)n_{(i)} = r_i M_i$$

and

$$n_i - \sum_{s=0}^{n_i-1} |\varphi_i^s(x_i)|^2 = \begin{cases} 0 & \text{if } d_i^{(n_i)} = 1 \\ \frac{r_i}{d_i^{(n_i)} - 1} (|\varphi_i^{n_i}(x_i)|^2 - 1) & \text{if } d_i^{(n_i)} > 1 \end{cases}$$

for all  $B < i \leq A$  and  $x_i \in G_i$ .

## Lemma

Let  $n \in \mathbf{P}$ .  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii\*).

$$D_n = n$$

**Property (ii\*)**  $B = A$  or there exist  $r_i < d_i^{(n_i)}$  nonnegative integers such that

$$(d_i^{(n_i)} - 1)n_{(i)} = r_i M_i$$

and

$$n_i - \sum_{s=0}^{n_i-1} |\varphi_i^s(x_i)|^2 = \begin{cases} 0 & \text{if } d_i^{(n_i)} = 1 \\ \frac{r_i}{d_i^{(n_i)} - 1} (|\varphi_i^{n_i}(x_i)|^2 - 1) & \text{if } d_i^{(n_i)} > 1 \end{cases}$$

for all  $B < i \leq A$  and  $x_i \in G_i$ .

## Lemma

Let  $n \in \mathbf{P}$ .  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii\*).

$$D_n = n$$

## Theorem

*Let  $G$  the complete product of finite groups where the dimensions of the representations appeared in the finite groups do not exceed the value 2. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

## Theorem

*Let  $G$  the complete product of finite groups where the square of the modulus of all of systems  $\varphi$  can only take integer values. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

## Theorem

*Suppose we order all of the systems  $\varphi$  such that the positive values of the identity are at the beginning of the systems. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

$$D_n = n$$

## Theorem

*Let  $G$  the complete product of finite groups where the dimensions of the representations appeared in the finite groups do not exceed the value 2. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

## Theorem

*Let  $G$  the complete product of finite groups where the square of the modulus of all of systems  $\varphi$  can only take integer values. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

## Theorem

*Suppose we order all of the systems  $\varphi$  such that the positive values of the identity are at the beginning of the systems. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

$$D_n = n$$

### Theorem

*Let  $G$  the complete product of finite groups where the dimensions of the representations appeared in the finite groups do not exceed the value 2. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

### Theorem

*Let  $G$  the complete product of finite groups where the square of the modulus of all of systems  $\varphi$  can only take integer values. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

### Theorem

*Suppose we order all of the systems  $\varphi$  such that the positive values of the identity are at the beginning of the systems. Then  $D_n = n$  if and only if  $n$  satisfies properties (i) and (ii).*

$$D_n = M_{A+1}$$

The finite group  $G_k$  has **property (N)** if there is an  $x_k \in G_k$  such that  $\varphi_k^{m_k-1}(x_k) = 0$ .

Several finite groups  $G_k$  can have property (N), so denote by  $N$  the smallest index of them, so  $G_N$  is the first group having property (N).

In the commutative case the index  $N$  does not exist.

For all  $k \in \mathbf{N}$  we can find the smaller number  $r \in \{1, 2, \dots, m_k - 1\}$  such that there exists an  $x_k \in G_k$  for which

$$\sum_{s=0}^r |\varphi_k^s(x_k)|^2 = m_k \quad \text{and} \quad \varphi_k^r(x_k) \neq 0$$

and denote this number by  $k^*$ .

$$D_n = M_{A+1}$$

The finite group  $G_k$  has **property (N)** if there is an  $x_k \in G_k$  such that  $\varphi_k^{m_k-1}(x_k) = 0$ .

Several finite groups  $G_k$  can have property (N), so denote by  $N$  the smallest index of them, so  $G_N$  is the first group having property (N).

In the commutative case the index  $N$  does not exist.

For all  $k \in \mathbf{N}$  we can find the smaller number  $r \in \{1, 2, \dots, m_k - 1\}$  such that there exists an  $x_k \in G_k$  for which

$$\sum_{s=0}^r |\varphi_k^s(x_k)|^2 = m_k \quad \text{and} \quad \varphi_k^r(x_k) \neq 0$$

and denote this number by  $k^*$ .

$$D_n = M_{A+1}$$

The finite group  $G_k$  has **property (N)** if there is an  $x_k \in G_k$  such that  $\varphi_k^{m_k-1}(x_k) = 0$ .

Several finite groups  $G_k$  can have property (N), so denote by  $N$  the smallest index of them, so  $G_N$  is the first group having property (N).

In the commutative case the index  $N$  does not exist.

For all  $k \in \mathbf{N}$  we can find the smaller number  $r \in \{1, 2, \dots, m_k - 1\}$  such that there exists an  $x_k \in G_k$  for which

$$\sum_{s=0}^r |\varphi_k^s(x_k)|^2 = m_k \quad \text{and} \quad \varphi_k^r(x_k) \neq 0$$

and denote this number by  $k^*$ .

The finite group  $G_k$  has **property (N)** if there is an  $x_k \in G_k$  such that  $\varphi_k^{m_k-1}(x_k) = 0$ .

Several finite groups  $G_k$  can have property (N), so denote by  $N$  the smallest index of them, so  $G_N$  is the first group having property (N).

In the commutative case the index  $N$  does not exist.

For all  $k \in \mathbf{N}$  we can find the smaller number  $r \in \{1, 2, \dots, m_k - 1\}$  such that there exists an  $x_k \in G_k$  for which

$$\sum_{s=0}^r |\varphi_k^s(x_k)|^2 = m_k \quad \text{and} \quad \varphi_k^r(x_k) \neq 0$$

and denote this number by  $k^*$ .

$$D_n = M_{A+1}$$

## Theorem

Let  $A \in \mathbf{N}$  and  $M_A \leq n < M_{A+1}$ . If the index  $N$  exists then  $D_n = M_{A+1}$  if and only if

$$\sum_{k=N}^A k^* M_k < n < M_{A+1}$$

If the index  $N$  does not exist then  $D_n < M_{A+1}$ .


$$D_n = M_{A+1}$$

## Theorem

Let  $A \in \mathbf{N}$  and  $M_A \leq n < M_{A+1}$ . If the index  $N$  exists then  $D_n = M_{A+1}$  if and only if

$$\sum_{k=N}^A k^* M_k < n < M_{A+1}$$

If the index  $N$  does not exist then  $D_n < M_{A+1}$ .



$D_n$  ( $n \leq 6^4$ ) on the complete product of  $S_3$  with new order

$$\{\varphi^0, \varphi^1, \varphi^4, \varphi^2, \varphi^5, \varphi^3\}$$

$$D_n = O(n)$$

The quotients  $\frac{D_n}{n}$  are not always bounded. For instance denote by  $d_k$  the dimension corresponding to  $\varphi_k^1$  and suppose that  $\varphi_k^1(e_k) = \sqrt{d_k}$ . Thus

$$\frac{D_n}{n} = \frac{1 + d_k}{2} \quad \text{if } n = 2M_k$$

so it can tend to infinity if  $d_k \rightarrow \infty$ .

## Lemma

*Let  $G$  the complete product of finite groups  $G_k$ . The quotients  $\frac{D_n}{n}$  are bounded for all  $n \in \mathbf{P}$  if and only if the quotients*

$$\frac{\sum_{s=0}^{r-1} |\varphi_k^s(x_k)|^2}{r}$$

*are bounded for all  $k \in \mathbf{N}$ ,  $0 < r \leq m_k$  and  $x_k \in G_k$ .*

$$D_n = O(n)$$

The quotients  $\frac{D_n}{n}$  are not always bounded. For instance denote by  $d_k$  the dimension corresponding to  $\varphi_k^1$  and suppose that  $\varphi_k^1(e_k) = \sqrt{d_k}$ . Thus

$$\frac{D_n}{n} = \frac{1 + d_k}{2} \quad \text{if } n = 2M_k$$

so it can tend to infinity if  $d_k \rightarrow \infty$ .

## Lemma

*Let  $G$  the complete product of finite groups  $G_k$ . The quotients  $\frac{D_n}{n}$  are bounded for all  $n \in \mathbf{P}$  if and only if the quotients*

$$\frac{\sum_{s=0}^{r-1} |\varphi_k^s(x_k)|^2}{r}$$

*are bounded for all  $k \in \mathbf{N}$ ,  $0 < r \leq m_k$  and  $x_k \in G_k$ .*

## Lemma

Let  $G$  a bounded group and  $xy^{-1} \in G \setminus \{e\}$ . Denote by  $j$  the index for which  $y \in I_j(x) \setminus I_{j+1}(x)$ . Then there exists a  $c > 0$  such that

$$|D_n(x, y)| \leq c \min \left\{ n, \frac{|\psi_{n^{(j+1)}}(x) \overline{\psi_{n^{(j+1)}}}(y)|}{|xy^{-1}|} \right\}$$

## Theorem (Dini's Test)

Let  $G$  the complete product of  $\mathcal{Q}_2$  with the product of the system appeared in the Table. Let  $x \in G$ ,  $f \in L^1(G)$  and suppose the function

$$g(y) := \frac{f(y) - f(x)}{|xy^{-1}|} \quad (y \neq x)$$

is integrable. Then  $S_n f(x) \rightarrow f(x)$  as  $n \rightarrow \infty$ .

## Lemma

Let  $G$  a bounded group and  $xy^{-1} \in G \setminus \{e\}$ . Denote by  $j$  the index for which  $y \in I_j(x) \setminus I_{j+1}(x)$ . Then there exists a  $c > 0$  such that

$$|D_n(x, y)| \leq c \min \left\{ n, \frac{|\psi_{n^{(j+1)}}(x) \overline{\psi_{n^{(j+1)}}}(y)|}{|xy^{-1}|} \right\}$$

## Theorem (Dini's Test)

Let  $G$  the complete product of  $\mathcal{Q}_2$  with the product of the system appeared in the Table. Let  $x \in G$ ,  $f \in L^1(G)$  and suppose the function

$$g(y) := \frac{f(y) - f(x)}{|xy^{-1}|} \quad (y \neq x)$$

is integrable. Then  $S_n f(x) \rightarrow f(x)$  as  $n \rightarrow \infty$ .